

8. Define simple harmonic motion in a straight line and show that

$$x = a \sin \omega t$$

can describe such motion, when  $x$  is the distance from a fixed point and  $a$ ,  $\omega$  and  $t$  have the usual meanings.

A particle  $p$ , of mass 5 kg, is connected by a light elastic string, of natural length 2 m and elastic constant 140 N/m to a fixed point  $q$  on a rough horizontal surface where the coefficient of friction is 1.  $p$  is released from rest at a point  $a$  where  $|qa| = 3$  m.

By considering the forces acting on  $p$  when its distance is  $(2.35 + x)$  m from  $q$ , prove that  $p$  moves in simple harmonic motion as long as the string remains taut. State the position of the centre,  $o$ , of the simple harmonic motion i.e.  $|qo|$  and write down the amplitude.

If the periodic time is assumed to be  $\frac{\pi}{\sqrt{7}}$  calculate the time taken by the particle to travel from  $a$  to a point 2 m from  $q$ .


Def<sup>n</sup> Notes.

$$x = a \sin \omega t \Rightarrow \frac{dx}{dt} = a\omega \cos \omega t \Rightarrow \frac{d^2x}{dt^2} = -a\omega^2 \sin \omega t$$


$$\Rightarrow \text{accel} = -\omega^2 (a \sin \omega t)$$

$$\Rightarrow \text{accel} = -\omega^2 x, \text{ qed.}$$

$a = \text{amplitude (case!)}$



Forces:



Accel =  $\frac{a}{\dots}$

Hooke's Law:

$$L = +140(2.35 + x - 2)$$

$$= +140(0.35 + x)$$

$$= 49 + 140x$$

Friction Law:

$$F = \mu R$$

$$\Rightarrow F = R, (\mu = 1)$$

W II:

$$\Sigma \vec{F} = m\vec{a}$$

in dir<sup>n</sup>

$$-L + F = 5a$$

$$-49 - 140x + R = 5a$$

$$-49 - 140x + 49 = 5a$$

$$\Rightarrow -140x = 5a$$

$$\boxed{-28x = a}$$

in dir<sup>n</sup>


$$R - W = 0$$

$$R - 5(9.8) = 0$$

$$\leftarrow R = 49$$


Particle is performing SHM with  $\omega = \sqrt{28}$  about a point 2.35 from wall.

Find Amplitude: Told  $v = 0$  where particle is 3m from wall.



$$v = 0, \text{ where } x = 0.65 \Rightarrow \boxed{A = 0.65}$$

Find time to go from a (Extreme) to point 2m from wall



Want time for particle to go from the + Extreme position to a position where  $x = -0.35$

$$x = A \cos \omega t \text{ (so start at extreme)}$$

At required position  $-0.35 = 0.65 \cos \sqrt{28}t$

$$x = -0.35 \Rightarrow -\frac{35}{65} = \cos \sqrt{28}t$$

$$\Rightarrow -0.5385 = \cos \sqrt{28}t \Rightarrow 2.139 = \sqrt{28}t$$

$$\Rightarrow 0.404 \text{ secs} = t$$